



## **OxyGeneo Facial Contraindications**

- ❖ A pacemaker or internal defibrillator, implanted neurostimulator, or other internal electric systems, metal implants or other implants in the treatment area, RF (not including dental implants and fillings)
- ❖ Impaired immune system due to immunosuppressive disease or use of immunosuppressive medications
- ❖ Use of depilatories, electrolysis, or waxing of the treated area in the past 7 days
- ❖ Sensory impairment in the treatment area. Diminished or exaggerated perception of the temperature changes. Conditions which could be adversely affected by heat
- ❖ Pregnant or nursing.
- ❖ Any active skin conditions within the applied area, such as sores, eczema, rash, fragile, swollen, burnt or injured skin, active acne, rosacea, dermatitis, psoriasis, or active Herpes Simplex. Numbness or loss of sensation
- ❖ Excessive fresh skin tan (within the last few days)
- ❖ Vascular disorders such as: telangiectasia, varicose veins, thrombosis, phlebitis in the applied area
- ❖ Severe concurrent disease such as: uncontrolled diabetes, epilepsy, lupus, nervous diseases, cardiac disorder. In such cases, consult your treating physician
- ❖ No Accutane usage for 12 months before treatment
- ❖ Deep chemical peels, laser skin resurfacing, or fractional RF surfacing within the last 3 months, Botox or Fillers within the last two weeks
- ❖ Known allergies to cosmetics or other products, or experienced severe allergic reactions like hives
- ❖ Use of active ingredients such as, but limited to Retinal, Hydroquinone, AHA's/BHA's, within 1 week of treatment
- ❖ Clients who are receiving chemotherapy/radiation/ have a history of cancer or have a deficient immune disorder, should consult a doctor - note will be required
- ❖ Active infection, cold sores, or Infectious disease, bacterial skin infections, fungal infections
- ❖ Grade IV acne, open lacerations in the area being treated
- ❖ History of skin disorders such as keloid scarring, abnormal wound healing, as well as very dry fragile skin
- ❖ Antibiotics, recent surgeries 8 weeks before treatment

If you are not sure about contraindications, please discuss at your consultation. Open communication is key to a successful treatment. Your safety will always be number 1